









**Scoring sheet of the SECONDS.** The scale features eight items (six mandatory - A, B, D, E, G, H and two conditional - C, F) presented in the order of administration. All required fields should be completed and the highest score used to determine the patient's behavioral diagnosis.

Patient : ..... Examiner : ..... Date : ..... Time : .....

**Simplified Evaluation of CONsciousness Disorders (SECONDS)**

<p>..... .....</p>	<p><b>A. Observation</b> </p>
<p>Command 1: ...../3 2: ...../3 3: ...../3 Written command: ...../3 → The patient responds at least twice for one of the commands (= score 6)</p>	<p><input type="checkbox"/> <b>B. Command-following</b> (score 6) 3 x 3 spoken commands 10'' interval between commands (1 x 3 written command if 0/3) Stop if 2 commands 3/3 </p>
<p style="writing-mode: vertical-rl; transform: rotate(180deg);">If command-following</p> <p>Code yes : ..... Code no : ..... Responses : .../5    o Verbal    o Autobiographical Correct : .../5    o Written    o Situational → The patient responds (even incorrectly) to at least 3 questions (= score 7) → The patient correctly responds to the 5 questions (= score 8)</p>	<p><input type="checkbox"/> <b>C. Communication</b> </p> <p><input type="checkbox"/> <b>Intentional</b> (score 7) <input type="checkbox"/> <b>Functional</b> (score 8) Autobiographical questions <i>Name (no), birth date (yes), name (yes), birth date (no), children (yes/no)</i> If incorrect answer(s): Situational questions <i>Place (yes), wearing a hat (no), place (no), touching hand (yes), touching face (no)</i></p>
<p>Horizontal : .../2                      Vertical : .../2 o Spontaneous   o Mirror            □ Manual eye-opening →The patient shows at least 2 visual pursuits of at least 2 seconds (= score 4)</p>	<p><input type="checkbox"/> <b>D. Visual pursuit</b> (score 4) </p> <p>Person/mirror, 30 cm from face Each movement on horizontal or vertical axes = 4''(→←↓↑)</p>
<p>Sup L : .../1                      Sup R : .../1 Inf L : .../1                      Inf R : .../1 o Spontaneous   o Mirror            □ Manual eye-opening →The patient shows at least 2 visual fixations of at least 2 seconds (= score 3)</p>	<p><input type="checkbox"/> <b>E. Visual fixation</b> (score 3) </p> <p>Person/mirror, 30 cm from face Present stimulus in each quadrant</p>
<p style="writing-mode: vertical-rl; transform: rotate(180deg);">If no command-following</p> <p>Localization: L : .../1    R : .../1 Anticipation: L : .../1    R : .../1 →The patient touches the point of stimulation at least once with the non-stimulated hand (= score 2) →The patient shows 2 anticipations (= score 6)</p>	<p><input type="checkbox"/> <b>F. Pain localization</b> (score 2) </p> <p>Inform patient 5'' pressure on nail bed 1 trial on each hand</p>
<p>..... Nb : ..... →The patient shows at least one oriented behavior (= score 5)</p>	<p><input type="checkbox"/> <b>G. Oriented behaviors</b> (score 5) </p> <p>E.g., scratching, grabbing sheets, holding bed, laughing or crying contextually,...</p>
<p>0-25% / 25-50% / 50-75% / 75-100% Spontaneously / Auditory / Tactile / Pain stimulations → The patient shows at least one eye-opening during the whole assessment (= score 1)</p>	<p><input type="checkbox"/> <b>H. Arousal</b> </p> <p><input type="checkbox"/> <b>Eye-opening</b> (score 1) <input type="checkbox"/> <b>No arousal</b> (score 0) Report the percentage of eye-opening time and administered stimulations</p>

**Diagnosis :** Coma (0) / UWS (1) / MCS- (2-5) / MCS+ (6-7) / EMCS (8)  
**Additional index point :** ... /100